



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Haemorrhagic bowel syndrome associated with ingestion of spoiled concentrate in two Brown Swiss cows

Braun, Ueli ; Gerspach, Christian ; Hässig, Michael

DOI: <https://doi.org/10.1136/vr.i4281>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-126320>

Journal Article

Accepted Version

Originally published at:

Braun, Ueli; Gerspach, Christian; Hässig, Michael (2016). Haemorrhagic bowel syndrome associated with ingestion of spoiled concentrate in two Brown Swiss cows. *Veterinary Record*, 179(6):152.

DOI: <https://doi.org/10.1136/vr.i4281>

Haemorrhagic bowel syndrome associated with ingestion of spoiled concentrate in two Brown Swiss cows

We wish to report an observation relating to the aetiology of haemorrhagic bowel syndrome (HBS). This is a serious disorder characterised by haemorrhage and secondary ileus of the small intestines in cattle. *Clostridium perfringens* type A (CPA) is thought to play an important role in the aetiology of this disorder (Elhafany et al. 2013) and was isolated from the faeces of 17 of 22 cows with HBS (Dennison et al. 2002); genotyping of ten of these samples yielded α -toxin-producing CPA (α -CPA) in five cows and α -CPA and β 2-toxin-producing CPA (β 2-CPA) in the remaining five. Another study found α -CPA in the intestinal contents of seven cows with HBS (Abutarbush and Radostits 2005), and analysis of tissue samples yielded α -CPA in three cows and β 2-CPA in four other cows with HBS (Ceci et al. 2006). Although there appears to be a causative link between CPA and HBS, experimental studies in which the proximal jejunum was inoculated with CPA failed to cause HBS in cows (Ewoldt and Anderson 2005). To our knowledge, studies on ingestion of feed contaminated with CPA have not been carried out, which prompted the present letter. Two Brown Swiss cows were recently referred to our clinic because of HBS that occurred after ingesting wet, clumped, foul-smelling, spoiled concentrate for several days. Analysis of the feed (miprolab GmbH, Göttingen, Germany) showed α -CPA, yeast ($8.32 \times 10^7/\text{g}$), fungi ($9.2 \times 10^6/\text{g}$) and non-specific bacterial flora. The timeline between ingestion of the spoiled feed and occurrence of HBS indicates that α -CPA, possibly in association with other degradation products of the feed, played a role in the aetiology of HBS in these cows.

Ueli Braun, Christian Gerspach, Michael Hässig, Department of Farm Animals, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 260, 8057 Zürich, Switzerland
e-mail: ueli.braun@uzh.ch

27

28 **References**

29 ABUTARBUSH, S. M. & RADOSTITS, O. M. (2005) Jejunal hemorrhage syndrome in dairy
30 and beef cattle: 11 cases (2001 to 2003). *Canadian Veterinary Journal* **46**, 711-715

31 CECI, L., PARADIES, P., SASANELLI, M., DE CAPRARIIS, D., GUARDA, F.,
32 CAPUCCHIO, M. T. & CARELLI, G. (2006) Haemorrhagic bowel syndrome in dairy
33 cattle: Possible role of *Clostridium perfringens* type A in the disease complex. *Journal of*
34 *Veterinary Medicine A* **53**, 518-523

35 DENNISON, A. C., VANMETRE, D. C., CALLAN, R. J., DINSMORE, P., MASON, G. L. &
36 ELLIS, R. P. (2002) Hemorrhagic bowel syndrome in dairy cattle: 22 cases (1997-2000).
37 *Journal of the American Veterinary Medical Association* **221**, 686-689

38 ELHAFANY, M. M, FRENCH, D. D. & BRAUN, U. (2013) Understanding jejunal
39 hemorrhage syndrome. *Journal of the American Veterinary Medical Association* **243**, 352-
40 358

41 EWOLDT, J. M. & ANDERSON, D. E. (2005) Determination of the effect of single
42 abomasal or jejunal inoculation of *Clostridium perfringens* type A in dairy cows. *Canadian*
43 *Veterinary Journal* **46**, 821-824